

Open Luttinger liquids

A. H. Castro Neto¹, C. de C. Chamon², and C. Nayak³

¹ *Dept. of Physics, University of California, Riverside, CA, 92521*

² *Dept. of Physics, University of Illinois at Urbana-Champaign, Urbana, IL, 61801*

³ *Institute for Theoretical Physics, University of California, Santa Barbara, CA, 93106*

(February 1, 2008)

We study the problem of Luttinger liquids interacting with an active environment. We are particularly interested in how dissipation affects the response and correlation functions of non-isolated Luttinger liquids. We show that the exchange of particles, energy, and momentum lead to changes in the exponents characterizing the various correlations functions. We discuss the importance of the zero mode physics in this context.

When a single quantum mechanical degree of freedom, X , is the cynosure in a system with many degrees of freedom, it is useful to regard the rest as a bath which damps the dynamics of X by dissipating its energy [1] and, potentially, by destroying its phase coherence [2]. Resistively shunted Josephson junctions and heavy particles in a metal are examples of physical systems which can be modelled by such a dissipative quantum mechanics [3]. It is very natural to wonder whether a similar treatment is applicable to a subsystem of reduced dimensionality which is spatially embedded in or at the boundary of a larger system which plays the role of the bath. Precisely this scenario appears to be realized in a number of different contexts.

Recent experiments in $\text{La}_{1.6-x}\text{Nd}_{0.4}\text{Sr}_x\text{CuO}_4$ and related compounds at $x = 1/8$ indicate a realization of a ‘striped’ phase in which Mott insulating antiferromagnetic regions are separated by anti-phase domain wall boundaries [4]. These domain walls, unlike those of $\text{La}_2\text{NiO}_{4+x}$, are not empty [4]. They have a filling fraction of $1/4$, or one electron for every two sites, so if the domain walls are treated as a system of quasi-one-dimensional wires running through the CuO_2 plane, they should be metallic – in fact Luttinger liquid – wires. However, the stripes are not simply Luttinger liquids because they interact with the antiferromagnetic spin fluctuations in the higher-dimensional Mott insulating regions [5], which can have several possible effects, including superconductivity [6].

Another example might be found in the quantum Hall regime [7]. Recall that the edge excitations of a fractional quantum Hall state ordinarily form a chiral Luttinger liquid [8]. However, these Luttinger liquids are no longer isolated when there is no gap in the bulk, as is the case at most even-denominator filling fractions [9] and in quantum Hall ferromagnets, which have gapless spin wave excitations in the limit of vanishing Zeeman energy [10]. Such effects of the bulk on the edge excitations could be detected in tunneling experiments [11].

One might worry that it does not always make sense to make this distinction between the subsystem and the bath when the subsystem has infinitely many degrees of freedom and interactions between the subsystem and the

rest of the system are not small. The excitations of the subsystem mix with those of the bath and the two cannot really be separated. Here, however, we consider the situation in which the subsystem is a spatially well-defined entity, as in the examples given above. In these cases, this division is particularly natural because certain types of transport measurements couple exclusively to the subsystem.

The preceding considerations serve as motivation. In this paper, we will be primarily concerned with setting up and solving a simple model of a Luttinger liquid interacting with a bath. First, we set up the model of a scalar field θ interacting with a bath. Integrating out the degrees of freedom of the bath induces non-local effective interactions in the scalar field. We focus on two novel aspects of the resulting theory: the non-local interactions which can be controlled by manipulating the bath, e.g. by varying the temperature; and the dissipation of the zero mode of the scalar field, as a result of which the conserved quantity (such as charge or spin) can leak into the bath along the whole length of the system (rather than just through external leads). We show how the combined effects modify the correlation functions of the operator $e^{i\theta}$. Finally, we discuss the implications for one-dimensional electron systems. We will only touch lightly on the applicability of this model to specific systems; a more thorough discussion, including comparisons with experimental data, will be reserved for a later publication.

The model we consider is a scalar field interacting with a bath of oscillators with some specifiable spectrum. Our primary concern is interacting one-dimensional electron systems, for which the scalar field is the bosonized representation of the electronic degrees of freedom. This representation is obtained by writing the fermion operator [12]

$$\psi(x, t) \propto \sum_{n \text{ odd}} e^{in\sqrt{\pi}(\theta(x, t) + k_F x)} e^{i\sqrt{\pi}\phi(x, t)} \quad (1)$$

in terms of non-independent boson fields ϕ and θ , which satisfy the commutation relation $[\phi(x), \theta(x')] = -i \text{sgn}(x - x')$. The action associated with these scalar

fields and their interaction with the bath (we use units such that $\hbar = k_B = 1$) is given by

$$S = S_{\mathcal{L}} + S_B + S_{\mathcal{L}B} \quad (2)$$

where $S_{\mathcal{L}}$ is the action of the scalar fields (which describes a Luttinger liquid),

$$S_{\mathcal{L}} = \frac{1}{2g} \int_{x,t} (\partial_\mu \theta)^2 = \frac{g}{2} \int_{x,t} (\partial_\mu \phi)^2, \quad (3)$$

where g is the Luttinger parameter that describes the forward scattering interactions between the fermions ($g < 1$, $g = 1$, $g > 1$ for repulsively interacting, free, and attractively interacting fermions, respectively) and the velocity of propagation is set to one ($\mu = 0, 1$ for the time and the one spatial coordinate). The fields ϕ and θ provide two dual descriptions of the Luttinger liquid. Alternatively, we may choose to work in a chiral basis, where we write $\theta = \chi_R + \chi_L$ and $\phi = g^{-1}(\chi_R - \chi_L)$. As a consequence of the equations of motion, the field ϕ can be expanded as

$$\chi_{R,L}(x,t) = \sqrt{\frac{g}{4\pi}} \left\{ X_{R,L} + P_{R,L} \frac{2\pi}{\ell} (t \mp x) + \sum_{n>0} \frac{i}{n} \left[a_{R,L;n} e^{-i\frac{2\pi}{\ell} n(t \mp x)} - c.c. \right] \right\} \quad (4)$$

where ℓ is the size of the system. The canonical quantization condition leads to the following commutation relations $[a_{R,L;n}, a_{R,L;m}^\dagger] = n\delta_{n,m}$, $[X_{R,L}, P_{R,L}] = i$. In the absence of the bath, $X_{R,L}, P_{R,L}, a_{R,L;n}, a_{R,L;n}^\dagger$ are time-independent. $P_{R,L}$ are right-moving and left-moving fermion numbers; $X_{R,L}$ are the conjugate phases. (In the language of conformal field theory, $P = P_R + P_L$ and $W = P_R - P_L$ are, respectively, the momentum and winding number of the scalar field.) When the coupling to the bath is turned on, these operators acquire time-dependence. Finally, the Hamiltonian associated with the decomposition reads,

$$H_{\mathcal{L}} = \frac{2\pi}{\ell} \left\{ \frac{P_R^2 + P_L^2}{2} + \sum_{n>0} \left[a_{R,n}^\dagger a_{R,n} + a_{L,n}^\dagger a_{L,n} \right] \right\}. \quad (5)$$

The dissipation arises from the interaction of the one dimensional system with the environment (the higher dimensional embedding system). This coupling will depend on the nature of the bath, and which degrees of freedom of the one dimensional system are sensitive to the fluctuations of the bath. More precisely, the environment could be a bath of photons, phonons, spin waves, or most generally, any set of collective excitations, which share the common feature of being described as a collection of harmonic oscillators. It is the physical nature of the bath that determines which sort of densities in the one dimensional system are affected by the environment, for example charge currents for coupling to a photon bath, and spin currents for coupling to a magnetic bath.

Here we will work with interactions which can be written in two ways: (A) In terms of either a coupling between a vector current and a vector field,

$$S_{\mathcal{L}B} = \int_{x,t} j_\mu(t,x) A^\mu(t,x, \mathbf{x}_\perp = 0), \quad (6)$$

where $j^\mu = \epsilon^{\mu\nu} \partial_\nu \theta / \sqrt{\pi}$ is the current obtained from the scalar field θ , and A^μ is a quantized harmonic field, and can describe any vector-like collective excitation for the bath; (B) In terms of density and a scalar field,

$$S_{\mathcal{L}B} = \int_{x,t} \partial_x \theta \sum_\alpha \lambda_\alpha \Phi_\alpha(t,x, \mathbf{x}_\perp = 0), \quad (7)$$

where α labels different modes, which couple to the density fluctuations $\delta\rho = \partial_x \theta / \sqrt{\pi}$ with different weights λ_α . In both cases the bath Hamiltonian is simply that of a set of harmonic oscillators with characteristic frequencies $\omega_{\mathbf{k},\alpha}$ which contain the information about the bath. Notice that the bath have extra degrees of freedom as compared to the 1-D system, which can be labelled by the perpendicular momenta $\mathbf{k}_\perp$ (the coupling to the 1-D system takes place at $\mathbf{x}_\perp = 0$).

This model is the natural generalization of that of a point particle interacting with a dissipative environment. The bath Hamiltonian is identical and the bilinear coupling of the bath to the particle's position has become a similar coupling to the field, θ or ϕ . The point particle is replaced by a free scalar field, which can be viewed, from a first-quantized standpoint, as a collection of non-interacting point particles. As in the point particle case, we can integrate out the bath to derive a non-local but *quadratic* effective action. This quadratic effective action, which we analyze below, is one of the main results of this paper.

Now, in contrast to the single particle, the field has many degrees of freedom labelled by the longitudinal momentum k . Each k mode of θ or ϕ is dissipated independently, so we can generalize the results for a single particle in a dissipative environment in a simple way. Integrating out the harmonic bath (*i.e.*, the fields A^μ or Φ_α), we are left with the following (Euclidean) action for the θ field:

$$S = \frac{1}{2} \int_{\omega,k} \left[\frac{1}{g} (\omega^2 + k^2) + J(k, \omega) \right] |\theta(k, \omega)|^2. \quad (8)$$

For each k mode, the properties of the bath can be summarized by a spectral function $J(k, \omega)$. Let us then turn to the analysis of the consequences of different forms for $J(k, \omega)$. Consider spectral functions that, for small k and ω behave as $J(k, \omega) \propto |k|^\alpha |\omega|^\beta$. We have the following possibilities: (i) $\alpha + \beta > 2$ - the effect of the bath on the long time or long distance physics is irrelevant. However, there may still be effects due to total charge dissipation if the zero mode couples to the bath. This will be considered later. (ii) $\alpha + \beta = 2$ - this is the marginal case.

The effect of $J(k, \omega)$ is to renormalize the Luttinger parameter g in front of the logarithmic correlation function for θ [13]. This occurs, for instance, in the case of a Luttinger liquid interacting with phonons, analyzed in [14]. The usual case of short range interactions, which shifts g , can be obtained by integrating out a bath of massive photons; the effect of the bath, as presented in this paper, is just a more general way to generate and think about interactions. We call attention to the fact that, in this picture, g arises from the bath and therefore may acquire dependence on an external parameter (*e.g.* temperature) affecting such bath. (iii) $\alpha + \beta < 2$ - in this case the bath leads to an energy gap in the spectrum of θ . One may worry that this will lead to inconsistencies if we look at correlations of $\psi \propto e^{i\theta}$, since the gap implies that $\lim_{t \rightarrow \infty} \langle \theta(t, 0) \theta(0, 0) \rangle = 0$, and so ψ acquires long range order. Kogut and Susskind [15] have studied this problem in the context of quark confinement. They have termed the vanishing of $\lim_{t \rightarrow \infty} \langle \theta(t, 0) \theta(0, 0) \rangle = 0$ the “seizing” of the vacuum: the interactions generated by the $J(k, \omega)$ are *long ranged*, which leads to *true ordering* in our 1+1 dimensional system. Even though there is a broken symmetry (shifts in θ), there are no Goldstone bosons associated with this broken symmetry; the infinite range interactions imply a finite energy cost even for long wavelength fluctuations. This case is especially interesting. The connected part of the electron Green’s function $\langle \psi(t, 0) \psi^\dagger(0, 0) \rangle_c$ vanishes except at $t = 0$. In other words, electrons are confined, so that this corresponds to a novel type of insulator phase where conduction is suppressed because free carriers are confined. The gap to excitations in the charged sector is proportional to the coupling to the bath.

As we mentioned previously, the effect of the bath is a more general way to think about interactions. Indeed, the bath can generate interactions which can be tuned via an external parameter, such as temperature or magnetic field, in such a way that one may switch among the three cases (i), (ii), (iii) above. Consider, for example, a bath of bosonic excitations in 2+1 dimensions, generating a $J(k, \omega) \propto k^2 / \sqrt{k^2 + m^2}$, where m is a mass gap. This mass measures a correlation length in the bath, $m \sim \xi^{-1}$, which can be, say, temperature dependent. Clearly, when $\xi \rightarrow \infty$, we are in case (iii). For any finite ξ , $J(k, \omega) \propto \xi k^2$, we are in case (ii) and the Luttinger parameter inherits the temperature dependence of the bath through ξ . For the cases (i) and (ii) corrections due to dissipation of the charge (zero mode) could be qualitatively important. Let us now turn to this problem.

If the subsystem exchanges charge (or spin) with the bath, in addition to energy and momentum, we must include the effects of the dissipation of the total charge of the subsystem. Here, we have in mind the possibility of charge being exchanged with the bath along the length of the system rather than a coupling to leads or point-contacts. To do so, we take proper care of the zero mode or total charge mode. We then allow for spectral functions that have a finite weight at $k = 0$:

$$J(k, \omega) = J_0(\omega) \delta(k) + J_{reg}(k, \omega) . \quad (9)$$

Several comments are in order. There are alternative ways of incorporating charge non-conservation in our model, such as coupling the bath directly to the operator $e^{i\theta}$. We have chosen a simple (and tractable) way of incorporating this phenomenon. As we show below, even the simple coupling (9) has interesting consequences. It is tempting to speculate on the possible similarity between the behavior resulting from different charge-dissipative couplings [16]. A second important point regarding (9) is the δ -function strength of the dissipation of the zero mode (which we justify in another way below), which is necessary to give a non-vanishing effect in the infinite-size limit. This, in turn, guarantees that the zero-mode dynamics has a non-vanishing effect on the Green function whereas, ordinarily, a single k mode with unit normalization would have vanishing contribution.

The zero mode of the scalar boson field θ can be read from the mode decomposition Eq. (4), $\theta_0(x, t) = \sqrt{\frac{\ell}{\pi}} \{X(t) + \pi W x / \ell\}$, where $X = (X_R + X_L)/2$, $P = P_R + P_L$, $W = P_L - P_R$, and in the last step we moved to the Heisenberg representation of X , P and W (notice that $X(t) = X + \pi P t / \ell$). Then $\theta(k = 0, t) = \sqrt{g/\pi} \ell X(t)$. We can also find a similar representation in terms of $\Theta = (X_R - X_L)/2$ which is the conjugate of the winding number W . Dissipation of the total current of the system enters through $X(t)$, whereas dissipation of the total charge enters through $\Theta(t)$. These two cases can be treated similarly, so for simplicity we consider below dissipation in just one, say X . Neglecting $J_{reg}(k, \omega)$ for the moment, we find an action for X of the form

$$S_X = \frac{1}{2} \int_{\omega} \left(\frac{\ell}{\pi} \omega^2 + \frac{g\ell^2}{2\pi^2} J_0(\omega) \right) |X(\omega)|^2 . \quad (10)$$

Notice that this is simply the dissipative action for a single particle of mass $M = \ell/\pi$, which here is our zero mode. The case of greatest interest, ohmic dissipation, $J_0(\omega) \propto |\omega|$ has been studied in great detail in the literature [3].

Since the dissipation is related to the collective mode of the bosonic field one expects to obtain Lagrangian (10) from a first-quantized viewpoint by treating the center of mass, $R_{CM}(t) = \sum_i x_i(t)/N$, of the bosons as the dissipative particle (N is the boson number). For ohmic dissipation the effective action (after the trace over the modes) reads [1]

$$S_d = \frac{\eta_P}{2\pi} \int dt dt' \frac{R_{CM}(t) R_{CM}(t')}{(t - t')^2} \quad (11)$$

where η_P is the friction coefficient. Using the definition of the center of mass and writing the action in terms of the fermion densities, $\rho(x, t) = \sum_i \delta(x - x_i(t)) = \rho_0 + \partial_x \theta(x, t) / \sqrt{\pi}$ where $\rho_0 = N/\ell$ is the average density, we find,

$$J_0(\omega) = \frac{\eta_P}{\ell^2 \rho_0^2} |\omega| \quad (12)$$

which gives the friction coefficient for the bosonic collective mode, η_P/ρ_0^2 .

The damping of the zero mode (12) leads to the dissipation into the bath of the conserved charge associated with the scalar field – either the charge or the spin of the bosonized electrons. This has profound consequences for the correlation functions of the operator $e^{i\theta}$. The $|\omega|$ dissipation term dominates the effective action (10) for low energies. Thus, for long times ($1 \gg t/\ell \gg 2\pi\rho_0^2/(g\eta_P)$), $\langle X(t)X(0) \rangle_c \approx \pi\rho_0^2/(g\eta_P) \ln[g\eta_P t/(2\pi\rho_0^2\ell)]$. In this case the correlation functions such as $\langle e^{i\theta(t,0)} e^{-i\theta(0,0)} \rangle$ are modified from their undamped forms by a factor of $t^{-\rho_0^2/\eta_P}$. Similarly, dissipation of charge would modify $\langle e^{i\phi(t,0)} e^{-i\phi(0,0)} \rangle$ by a factor $t^{-\rho_0^2/\eta_W}$, where η_W is the friction coefficient for the total charge. These corrections explicitly alter the electron propagator in the 1-D system:

$$G(x, t) = \langle \psi(x, t) \psi^\dagger(0, 0) \rangle \approx \frac{G_0(x, t)}{\left[\frac{g\eta_P t}{2\pi\rho_0^2\ell} \right]^{\pi\rho_0^2/\eta_P} \left[\frac{\eta_W t}{2\pi g\rho_0^2\ell} \right]^{\pi\rho_0^2/\eta_W}} \quad (13)$$

where $G_0(x, t)$ is the correlation function in the absence of charge dissipation. Our conclusion is therefore that for long times the electron correlation function decays like $t^{-\gamma}$, where $\gamma = (g + g^{-1})/2 + \pi\rho_0^2(\eta_P^{-1} + \eta_W^{-1})$. This correlation function decays faster than the usual Luttinger liquid correlation function. In many examples of interest, $\eta_{P,W}$ will be temperature-dependent, and the Green function will have unusual temperature dependence.

This is precisely the dissipation of charge that we mentioned earlier. As we will discuss further, the damping changes the Luttinger liquid exponents for tunneling at a point contact. Naively, the physics behind η is quite different from the physics relating to the Luttinger liquid parameter, g . The former reflects the diffusion of charge away from the Luttinger liquid (in a position independent way); the latter reflects the reduced density of states at the Fermi points due to repulsive interactions. However, for point contact tunneling, they have a similar effect because the Luttinger liquid plays the role of a bath and, hence, $\frac{1}{2}(g + g^{-1})$ and $\pi\rho_0^2(\eta_P^{-1} + \eta_W^{-1})$ play similar roles.

In an electron system which is not spin-polarized, there is a non-trivial spin sector. The charge sector is similar to that of spinless (i.e. spin-polarized) electrons; if it is coupled to a bath, the charge dynamics is determined by the previous results. If, on the other hand, the spin sector is coupled to a bath while the charge sector is undamped, both charge and spin dynamics can be anomalous. The latter is not surprising; the former occurs because electron tunneling necessarily involves the recombination of spinon and holons. In the abelian bosonization scheme, both the charge and spin sectors are described by free scalar fields with Luttinger liquid parameters g_c and $g_s = 2$. The electron operator, $e^{i\phi_c} e^{i\phi_s}$ is a dimension $(1/g_c + 1/g_s)/2$ operator. In the absence of dissipation, this determines the temperature dependence of

tunneling at a point contact, $G \sim T^{2(1-1/g_c-1/g_s)}$. In the presence of dissipation, the dimension of the electron tunneling operator is shifted by $\pi\rho_0^2\eta_{s,c}^{-1}$ through the shift in the spinon operator, so $G \sim T^{2(1-1/g_c-1/g_s-\pi\rho_0^2\eta_{s,c}^{-1})}$, in complete analogy with the previous discussion.

In summary, we have, in this paper, considered the simplest possible ‘toy model’ of a Luttinger liquid interacting with a dissipative bath. Remarkably, we find interesting behavior even in this model such as modified Luttinger liquid exponents which depend on the spectrum of the bath. Possible consequences for the QHE and cuprates will be considered elsewhere.

We thank A. Balatsky, G. Castilla, E. Fradkin, D. E. Freed, M. R. Geller, A.W.W. Ludwig, S. Kivelson, C. Mudry, S. H. Simon, and X.-G. Wen for discussions. The authors acknowledge the Aspen Center for Physics for the hospitality during the initial stages of this work. This work was supported in part by the National Science Foundation through grants NSF-DMR-94-24511, NSF-DMR-89-20538 and NSF-ITP-97-029.

-
- [1] A.O.Caldeira and A.J.Leggett, *Annals of Physics* **149**, 374 (1983); *Physica* **121 A**, 587 (1983).
 - [2] A.J.Leggett *et al.*, *Rev.Mod.Phys.* **59**, 1 (1987).
 - [3] A. Schmid, *Phys.Rev.Lett.* **51**, 1506 (1983); F.Guinea *et al.*, *Phys.Rev.Lett.* **54**, 263 (1985); S.Chakravarty *et al.*, *Phys.Rev.Lett.* **56**, 2303 (1986); Yu. Kagan and N.V.Prokofev, in *Quantum tunneling in condensed media*, Yu. Kagan and A.J.Leggett Eds. (Elsevier, 1992), pg. 37.
 - [4] Tranquada *et al.*, *Nature*, **375**, 561, (1995); Tranquada *et al.*, *Phys.Rev.B* **54**, 7489 (1996).
 - [5] C. Nayak, and F. Wilczek, *Int. J. Mod. Phys. B* **10**, 2125 (1996); J. Zaanen *et al.*, *Phys. Rev. B* **53**, 8671 (1996); A. H. Castro Neto, cond-mat 9611146.
 - [6] V. J. Emery *et al.*, cond-mat 9610094;
 - [7] R.E. Prange and S.M. Girvin eds., *The Quantum Hall Effect*, 2nd ed. (Springer-Verlag, New York, 1990).
 - [8] X. G. Wen, *Phys. Rev. B* **41**, 12838 (1990).
 - [9] B. I. Halperin, P. A. Lee, and N. Read, *Phys. Rev. B* **47**, 7312 (1993).
 - [10] S. Sondhi, *et al.*, *Phys. Rev. B* **47**, 16419 (1993).
 - [11] A. M. Chang, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* **77**, 2538 (1996).
 - [12] F. D. M. Haldane, *J. Phys. C* **14**, 2585 (1981); *Phys. Rev. Lett.* **47**, 1840 (1981).
 - [13] Even when $\alpha \neq \beta$, the spatial and temporal dependence of the correlation functions are affected in the same way so long as $\alpha + \beta = 2$.
 - [14] T. Martin and D. Loss, *Int. J. Mod. Phys. B*, **9**, 495 (1995). O. Heinonen and S. Eggert, *Phys. Rev. Lett* **77**, 358 (1996).
 - [15] J.Kogut and L.Susskind, *Phys.Rev.D* **11**, 3594 (1975).
 - [16] In a recent preprint, cond-mat/9703246, Shytov, Levitov,

and Halperin have treated the equilibration between edge channels in a way which is reminiscent of ours, namely by assuming that the charge transfer can be described by a quadratic action, rather than a complicated action with exponential operators. As in this paper, the non-trivial behavior results from the subsequent exponentiation of these quadratic fields.